

wherein Z is H, OH, YNOX, OR<sub>5</sub>, -NR<sub>5</sub>R<sub>6</sub> and R<sub>1</sub> and R<sub>2</sub>, are as defined in claim 1; and R<sub>7</sub> is cycloalkyl of 3-6 carbon atoms; 5-10 membered cycloheteroalkyl; alkyl of 1-18 carbon atoms; alkenyl of 2-18 carbon atoms having from 1 to 3 double bonds; alkynyl of 2-18 carbon atoms having from 1 to 3 triple bonds; or -SiR<sub>8</sub>R<sub>9</sub>R<sub>10</sub>;

R<sub>8</sub>, R<sub>9</sub>, and R<sub>10</sub> are each, independently, aryl; 4-8 membered heteroaryl having 1-3 heteroatoms selected from N, NR<sub>4</sub>, O and S; cycloalkyl of 3-6 carbon atoms; 5-10 membered cycloheteroalkyl; alkyl of 1-18 carbon atoms; alkenyl of 2-18 carbon atoms having from 1 to 3 double bonds; alkynyl of 2-18 carbon atoms having from 1 to 3 triple bonds; or two of R<sub>8</sub>, R<sub>9</sub>, and R<sub>10</sub> taken together with the silicon atom to which they are attached form a heterocyclic ring of 5 or 6 members;

in the presence of a Lewis acid or fluoride reagent in an ether organic solvent at temperatures ranging from about -78°C to about 30°C to produce an alpha-sulfonyl carbonyl compound of formula V; any reactive substituent group(s) being protected during the reaction and removed thereafter; and further if desired isolating any chiral or stereoisomeric product as an individual isomer.

#### REMARKS

The present invention relates to certain alpha-sulfonyl hydroxamic acid derivatives, pharmaceutical composition containing them and a method of preparation.

Claims 1-52 have been rejected under 35 USC 102(a) as being anticipated by WO 00/71514 ("Barta"). The Examiner contends that Barta teaches the claimed sulfonyl hydroxamic acid derivatives.

Applicants respectfully traverse the rejection. Barta is concerned with beta-sulfonyl hydroxamic acid derivatives whereas the present invention relates to alpha-sulfonyl hydroxamic acid derivatives. Applicants have reviewed the passages cited by the Examiner, but do not believe that Barta anticipates the present invention.

Claims 1-44 have been rejected under 35 USC 102(b) as being anticipated by J. Org. Chem. Vol. 55, pp. 1125-1126 ("Kende").


Applicants again respectfully traverse the rejection. Although Kende is concerned with alpha-sulfonyl hydroxamic acid derivatives. Kende uses only alkyl and cycloalkyl esters. Applicants have amended the claims such that R<sub>1</sub> and R<sub>2</sub> are aryl, heteroaryl or cycloheteroalkyl. Applicants believe that the claims as amended are patentable over Kende.

Claims 1-52 have been rejected under 35 USC 112, first paragraph, because the Examiner contends that the specification while being enabling for R<sub>1</sub>, R<sub>2</sub> piperidine derivative does not provide enablement for heterocyclic containing derivatives, particularly thiomorpholine derivatives.

Applicants respectfully traverse the rejection. While the Examiner is correct that Schemes I, II, and III all show the preferred piperidine group, the specification clearly indicates that these are preferred routes (see paragraph bridging pages 23 and 24). Applicants believe that one of ordinary skill in the art after reading the specification (including the schemes and examples) would be able to substitute any claimed group for the clearly depicted piperidine group without undue experimentation. Specifically in response to the Examiner's stated concern with thiomorpholides, one of ordinary skill in the art would be able to make the compounds using the process as claimed.

Attached hereto is a marked-up version of the changes made to the specification and claims by the current amendment. The attached page is captioned **"Version with Markings to Show Changes Made."**

Applicants believe that the present application is in condition for allowance and respectfully request that the Examiner enter the amendment, reconsider the rejections in light of the remarks herein and allow the application. Favorable treatment of the application is earnestly solicited.

  
John W. Hogan, Jr.

Reg. No. 32,703


Wyeth  
Patent Law Department  
Five Giralda Farms  
Madison, NJ 07940  
Tel. No. (973) 683-2152

**Version with Markings to Show Changes Made**

**In the Claims**

Claim 1 has been amended as follows:

1. (Amended) A method of preparing alpha-sulfonyl derivatives of the formula V:



wherein Z is H, OH, -NYOX, -OR<sub>5</sub> or -NR<sub>5</sub>R<sub>6</sub>;

X is hydrogen, alkyl of 1-6 carbon atoms, benzyl, hydroxyethyl, t-butyldimethylsilyl, trimethylsilyl or tetrahydropyranyl;

Y is hydrogen, alkyl of 1-6 carbon atoms, aryl of 6 to 10 carbon atoms, 5-10 membered heteroaryl having 1-3 heteroatoms selected from N, NR<sub>4</sub>, O and S, cycloalkyl of 3-6 carbon atoms, 5-10 membered cycloheteroalkyl; wherein said alkyl, aryl, heteroaryl, cycloalkyl and cycloheteroalkyl group of Y is optionally substituted on any atom capable of substitution, with 1 to 3 substituents selected from the group consisting of halogen, alkyl of 1-6 carbon atoms; alkenyl of 2-6 carbon atoms having from 1 to 3 double bonds; alkynyl of 2-6 carbon atoms having from 1 to 3 triple bonds, cycloalkyl of 3-6 carbon atoms, -OR<sub>5</sub>, =O, -CN, -COR<sub>5</sub>, perfluoroalkyl of 1-4 carbon atoms, -O-perfluoroalkyl of 1-4 carbon atoms, -CONR<sub>5</sub>R<sub>6</sub>, -S(O)<sub>n</sub>R<sub>5</sub>, -OPO(OR<sub>5</sub>)OR<sub>6</sub>, -PO(OR<sub>5</sub>)R<sub>6</sub>, -OC(O)OR<sub>5</sub>, -OR<sub>5</sub>NR<sub>5</sub>R<sub>6</sub>, -OC(O)NR<sub>5</sub>R<sub>6</sub>, -C(O)NR<sub>5</sub>OR<sub>6</sub>, -COOR<sub>5</sub>, -SO<sub>3</sub>H, -NR<sub>5</sub>R<sub>6</sub>, -N[(CH<sub>2</sub>)<sub>2</sub>]<sub>2</sub>NR<sub>5</sub>, -NR<sub>5</sub>COR<sub>6</sub>, -NR<sub>5</sub>COOR<sub>6</sub>, SO<sub>2</sub>NR<sub>5</sub>R<sub>6</sub>, -NO<sub>2</sub>, -N(R<sub>5</sub>)SO<sub>2</sub>R<sub>6</sub>, -NR<sub>5</sub>CONR<sub>5</sub>R<sub>6</sub>, -NR<sub>5</sub>C(=NR<sub>6</sub>)NR<sub>5</sub>R<sub>6</sub>, -NR<sub>5</sub>C(=NR<sub>6</sub>)N(SO<sub>2</sub>R<sub>5</sub>)R<sub>6</sub>, -NR<sub>5</sub>C(=NR<sub>6</sub>)N(C=OR<sub>5</sub>)R<sub>6</sub>, -tetrazol-5-yl, -SO<sub>2</sub>NHCN, -SO<sub>2</sub>NHCONR<sub>5</sub>R<sub>6</sub>, phenyl, heteroaryl and 5-10 membered cycloheteroalkyl;

R<sub>1</sub> and R<sub>2</sub> are each, independently, ~~hydrogen~~; aryl of 6 to 10 carbon atoms; 5-10 membered heteroaryl having 1-3 heteroatoms selected from N, NR<sub>4</sub>, O and S; ~~cycloalkyl of 3-6 carbon atoms~~; 5-10 membered cycloheteroalkyl; ~~alkyl of 1-18 carbon atoms~~; ~~alkenyl of 2-18 carbon atoms~~

atoms having 1 to 3 double bonds; alkynyl of 2-18 carbon atoms having from 1 to 3 triple bonds; or R<sub>1</sub> and R<sub>2</sub> taken together with the carbon atom to which they are attached form a cycloalkyl ring of 3-8 carbon atoms or a 5-10 membered cycloheteroalkyl ring; and wherein the aryl, heteroaryl, ~~cycloalkyl~~, or cycloheteroalkyl, ~~alkyl~~, ~~alkenyl~~, and ~~alkynyl~~, may be optionally substituted on any atom capable of substitution with from 1 to 3 substituents selected from halogen, alkyl of 1-6 carbon atoms; alkenyl of 2-6 carbon atoms having from 1 to 3 double bonds; alkynyl of 2-6 carbon atoms having from 1 to 3 triple bonds, cycloalkyl of 3-6 carbon atoms, -OR<sub>5</sub>, =O, -CN, -COR<sub>5</sub>, perfluoroalkyl of 1-4 carbon atoms, -O-perfluoroalkyl of 1-4 carbon atoms, -CONR<sub>5</sub>R<sub>6</sub>, -S(O)<sub>n</sub>R<sub>5</sub>, -OPO(OR<sub>5</sub>)OR<sub>6</sub>, -PO(OR<sub>5</sub>)R<sub>6</sub>, -OC(O)OR<sub>5</sub>, -OR<sub>5</sub>NR<sub>5</sub>R<sub>6</sub>, -OC(O)NR<sub>5</sub>R<sub>6</sub>, -C(O)NR<sub>5</sub>OR<sub>6</sub>, -COOR<sub>5</sub>, -SO<sub>3</sub>H, -NR<sub>5</sub>R<sub>6</sub>, -N[(CH<sub>2</sub>)<sub>2</sub>]<sub>2</sub>NR<sub>5</sub>, -NR<sub>5</sub>COR<sub>6</sub>, -NR<sub>5</sub>COOR<sub>6</sub>, SO<sub>2</sub>NR<sub>5</sub>R<sub>6</sub>, -NO<sub>2</sub>, -N(R<sub>5</sub>)SO<sub>2</sub>R<sub>6</sub>, -NR<sub>5</sub>CONR<sub>5</sub>R<sub>6</sub>, -NR<sub>5</sub>C(=NR<sub>6</sub>)NR<sub>5</sub>R<sub>6</sub>, -NR<sub>5</sub>C(=NR<sub>6</sub>)N(SO<sub>2</sub>R<sub>5</sub>)R<sub>6</sub>, -NR<sub>5</sub>C(=NR<sub>6</sub>)N(C=OR<sub>5</sub>)R<sub>6</sub>, -tetrazol-5-yl, -SO<sub>2</sub>NHCN, -SO<sub>2</sub>NHCONR<sub>5</sub>R<sub>6</sub>, phenyl, heteroaryl and 5-10 membered cycloheteroalkyl;


R<sub>3</sub> is alkyl of 1-18 carbon atoms, alkenyl of 2-18 carbon atoms having 1 to 3 double bonds, alkynyl of 2-18 carbon atoms having from 1 to 3 triple bonds, cycloalkyl of 3-6 carbon atoms, 5-10 membered cycloheteroalkyl, aryl of 6 to 10 carbon atoms, 5-6 membered heteroaryl having 1-3 heteroatoms selected from N, NR<sub>4</sub>, O, and S; wherein said alkyl, alkenyl, alkynyl, cycloalkyl, cycloheteroalkyl, aryl and heteroaryl of R<sub>3</sub> may optionally be substituted on any atom capable of substitution with from 1 to 3 substituents selected from halogen, alkyl of 1-6 carbon atoms; alkenyl of 2-6 carbon atoms having from 1 to 3 double bonds; alkynyl of 2-6 carbon atoms having from 1 to 3 triple bonds, cycloalkyl of 3-6 carbon atoms, -OR<sub>5</sub>, =O, -CN, -COR<sub>5</sub>, perfluoroalkyl of 1-4 carbon atoms, -O-perfluoroalkyl of 1-4 carbon atoms, -CONR<sub>5</sub>R<sub>6</sub>, -S(O)<sub>n</sub>R<sub>5</sub>, -OPO(OR<sub>5</sub>)OR<sub>6</sub>, -PO(OR<sub>5</sub>)R<sub>6</sub>, -OC(O)OR<sub>5</sub>, -OR<sub>5</sub>NR<sub>5</sub>R<sub>6</sub>, -OC(O)NR<sub>5</sub>R<sub>6</sub>, -C(O)NR<sub>5</sub>OR<sub>6</sub>, -COOR<sub>5</sub>, -SO<sub>3</sub>H, -NR<sub>5</sub>R<sub>6</sub>, -N[(CH<sub>2</sub>)<sub>2</sub>]<sub>2</sub>NR<sub>5</sub>, -NR<sub>5</sub>COR<sub>6</sub>, -NR<sub>5</sub>COOR<sub>6</sub>, SO<sub>2</sub>NR<sub>5</sub>R<sub>6</sub>, -NO<sub>2</sub>, -N(R<sub>5</sub>)SO<sub>2</sub>R<sub>6</sub>, -NR<sub>5</sub>CONR<sub>5</sub>R<sub>6</sub>, -NR<sub>5</sub>C(=NR<sub>6</sub>)NR<sub>5</sub>R<sub>6</sub>, -NR<sub>5</sub>C(=NR<sub>6</sub>)N(SO<sub>2</sub>R<sub>5</sub>)R<sub>6</sub>, -NR<sub>5</sub>C(=NR<sub>6</sub>)N(C=OR<sub>5</sub>)R<sub>6</sub>, -tetrazol-5-yl, -SO<sub>2</sub>NHCN, -SO<sub>2</sub>NHCONR<sub>5</sub>R<sub>6</sub>, phenyl, heteroaryl and 5-10 membered cycloheteroalkyl;

R<sub>4</sub> is hydrogen; aryl; aralkyl, heteroaryl; heteroaralkyl, alkyl of 1-6 carbon atoms; cycloalkyl of 3-6 carbon atoms; -C(O)<sub>n</sub>R<sub>5</sub>, -CONR<sub>5</sub>R<sub>6</sub> or SO<sub>2</sub>R<sub>5</sub>;

R<sub>5</sub> and R<sub>6</sub> are each independently hydrogen, optionally substituted aryl; 4-8 membered heteroaryl having 1-3 heteroatoms selected from N, NR<sub>4</sub>, O and S; cycloalkyl of 3-6 carbon atoms; 5-10 membered cycloheteroalkyl; alkyl of 1-18 carbon atoms; alkenyl of 2-18 carbon atoms or alkynyl of 2-18 carbon atoms; or R<sub>5</sub> and R<sub>6</sub> taken together with the nitrogen atom to which they are attached may form a 5-10 membered cycloheteroalkyl ring; and


n is 1 or 2; or a pharmaceutical salt thereof,

which comprises reacting a sulfonyl fluoride of the formula III



III

wherein R<sub>3</sub>' is as hereinabove defined for R<sub>3</sub> with the proviso that R<sub>3</sub>' does not contain a group that can form an anion under basic conditions; with a carbonyl compound of the formula IV:




IV

wherein Z is H, OH, YNOX, -NR<sub>5</sub>R<sub>6</sub> or OR<sub>5</sub>, and X, Y, R<sub>1</sub>, R<sub>2</sub>, R<sub>5</sub>, and R<sub>6</sub> are as hereinabove defined; in the presence of a metal hydride or amide base in an ether organic solvent at temperatures from about -78°C to about 30°C to produce an alpha-sulfonyl carbonyl compound of formula V;

any reactive substituent group(s) being protected during the reaction and removed thereafter; and further if desired isolating any chiral or stereoisomeric product as an individual isomer.

Claim 15 has been amended as follows:

15. (Amended) A method of preparing alpha-sulfonyl derivatives of the formula V:



wherein Z is H, OH, -NYOX, -OR<sub>5</sub> or -NR<sub>5</sub>R<sub>6</sub>;


R<sub>1</sub> and R<sub>2</sub> are each, independently, ~~hydrogen~~; aryl of 6 to 10 carbon atoms; 5-10 membered heteroaryl having 1-3 heteroatoms selected from N, NR<sub>4</sub>, O and S; ~~cycloalkyl of 3-6 carbon atoms; 5-10 membered cycloheteroalkyl; alkyl of 1-18 carbon atoms; alkenyl of 2-18 carbon atoms having 1 to 3 double bonds; alkynyl of 2-18 carbon atoms having from 1 to 3 triple bonds~~; or R<sub>1</sub> and R<sub>2</sub> taken together with the carbon atom to which they are attached form a ~~cycloalkyl ring of 3-8 carbon atoms or a 5-10 membered cycloheteroalkyl ring~~; and wherein the aryl, heteroaryl, ~~cycloalkyl~~, or cycloheteroalkyl, alkyl, alkenyl, and alkynyl, may be optionally substituted on any atom capable of substitution with from 1 to 3 substituents selected from halogen, alkyl of 1-6 carbon atoms; alkenyl of 2-6 carbon atoms having from 1 to 3 double bonds; alkynyl of 2-6 carbon atoms having from 1 to 3 triple bonds, cycloalkyl of 3-6 carbon atoms, -OR<sub>5</sub>, =O, -CN, -COR<sub>5</sub>, perfluoroalkyl of 1-4 carbon atoms, -O-perfluoroalkyl of 1-4 carbon atoms, -CONR<sub>5</sub>R<sub>6</sub>, -S(O)<sub>n</sub>R<sub>5</sub>, -OPO(OR<sub>5</sub>)OR<sub>6</sub>, -PO(OR<sub>5</sub>)R<sub>6</sub>, -OC(O)OR<sub>5</sub>, -OR<sub>5</sub>NR<sub>5</sub>R<sub>6</sub>, -OC(O)NR<sub>5</sub>R<sub>6</sub>, -C(O)NR<sub>5</sub>OR<sub>6</sub>, -COOR<sub>5</sub>, -SO<sub>3</sub>H, -NR<sub>5</sub>R<sub>6</sub>, -N[(CH<sub>2</sub>)<sub>2</sub>]<sub>2</sub>NR<sub>5</sub>, -NR<sub>5</sub>COR<sub>6</sub>, -NR<sub>5</sub>COOR<sub>6</sub>, SO<sub>2</sub>NR<sub>5</sub>R<sub>6</sub>, -NO<sub>2</sub>, -N(R<sub>5</sub>)SO<sub>2</sub>R<sub>6</sub>, -NR<sub>5</sub>CONR<sub>5</sub>R<sub>6</sub>, -NR<sub>5</sub>C(=NR<sub>6</sub>)NR<sub>5</sub>R<sub>6</sub>, -NR<sub>5</sub>C(=NR<sub>6</sub>)N(SO<sub>2</sub>R<sub>5</sub>)R<sub>6</sub>, -NR<sub>5</sub>C(=NR<sub>6</sub>)N(C=OR<sub>5</sub>)R<sub>6</sub>, -tetrazol-5-yl, -SO<sub>2</sub>NHCN, -SO<sub>2</sub>NHCONR<sub>5</sub>R<sub>6</sub>, phenyl, heteroaryl and 5-10 membered cycloheteroalkyl;

R<sub>3</sub>' is alkyl of 1-18 carbon atoms, alkenyl of 2-18 carbon atoms having 1 to 3 double bonds, alkynyl of 2-18 carbon atoms having from 1 to 3 triple bonds, cycloalkyl of 3-6 carbon atoms, 5-10 membered cycloheteroalkyl, aryl of 6 to 10 carbon atoms, 5-6 membered heteroaryl having 1-3 heteroatoms selected from N, NR<sub>4</sub>, O, and S; wherein said alkyl, alkenyl, alkynyl, cycloalkyl, cycloheteroalkyl, aryl and heteroaryl of R<sub>3</sub> may optionally be substituted on any atom capable of substitution with from 1 to 3 substituents selected from halogen, alkyl of 1-6 carbon atoms; alkenyl of 2-6 carbon atoms having from 1 to 3 double bonds; alkynyl of 2-6 carbon atoms having from 1 to 3 triple bonds, cycloalkyl of 3-6 carbon atoms, -OR<sub>5</sub>, =O, -CN, -COR<sub>5</sub>, perfluoroalkyl of 1-4 carbon atoms, -O-perfluoroalkyl of 1-4 carbon atoms, -CONR<sub>5</sub>R<sub>6</sub>, -S(O)<sub>n</sub>R<sub>5</sub>, -OPO(OR<sub>5</sub>)OR<sub>6</sub>, -PO(OR<sub>5</sub>)R<sub>6</sub>, -OC(O)OR<sub>5</sub>, -OR<sub>5</sub>NR<sub>5</sub>R<sub>6</sub>, -OC(O)NR<sub>5</sub>R<sub>6</sub>, -C(O)NR<sub>5</sub>OR<sub>6</sub>, -COOR<sub>5</sub>, -SO<sub>3</sub>H, -NR<sub>5</sub>R<sub>6</sub>, -N[(CH<sub>2</sub>)<sub>2</sub>]<sub>2</sub>NR<sub>5</sub>, -NR<sub>5</sub>COR<sub>6</sub>, -NR<sub>5</sub>COOR<sub>6</sub>, SO<sub>2</sub>NR<sub>5</sub>R<sub>6</sub>, -NO<sub>2</sub>, -N(R<sub>5</sub>)SO<sub>2</sub>R<sub>6</sub>, -NR<sub>5</sub>CONR<sub>5</sub>R<sub>6</sub>,


-NR<sub>5</sub>C(=NR<sub>6</sub>)NR<sub>5</sub>R<sub>6</sub>, -NR<sub>5</sub>C(=NR<sub>6</sub>)N(SO<sub>2</sub>R<sub>5</sub>)R<sub>6</sub>, -NR<sub>5</sub>C(=NR<sub>6</sub>)N(C=OR<sub>5</sub>)R<sub>6</sub>,  
-tetrazol-5-yl, -SO<sub>2</sub>NHCN, -SO<sub>2</sub>NHCONR<sub>5</sub>R<sub>6</sub>, phenyl, heteroaryl and 5-10 membered  
cycloheteroalkyl; provided that R<sub>3</sub>' does not contain a group that can form an anion under  
basic conditions;

or a pharmaceutically acceptable salt thereof, which comprises the steps of :

a) reacting a sulfonyl fluoride of formula III:



wherein R<sub>3</sub>' is as defined in claim 1; with an enol ether of formula VIII:



wherein Z is H, OH, YNOX, OR<sub>5</sub>, -NR<sub>5</sub>R<sub>6</sub> and R<sub>1</sub> and R<sub>2</sub>, are as defined in claim 1; and  
R<sub>7</sub> is cycloalkyl of 3-6 carbon atoms; 5-10 membered cycloheteroalkyl; alkyl of 1-18 carbon  
atoms; alkenyl of 2-18 carbon atoms having from 1 to 3 double bonds; alkynyl of 2-18 carbon  
atoms having from 1 to 3 triple bonds; or -SiR<sub>8</sub>R<sub>9</sub>R<sub>10</sub>;

R<sub>8</sub>, R<sub>9</sub>, and R<sub>10</sub> are each, independently, aryl; 4-8 membered heteroaryl having 1-3  
heteroatoms selected from N, NR<sub>4</sub>, O and S; cycloalkyl of 3-6 carbon atoms; 5-10 membered  
cycloheteroalkyl; alkyl of 1-18 carbon atoms; alkenyl of 2-18 carbon atoms having from 1 to  
3 double bonds; alkynyl of 2-18 carbon atoms having from 1 to 3 triple bonds; or two of R<sub>8</sub>,  
R<sub>9</sub>, and R<sub>10</sub> taken together with the silicon atom to which they are attached form a  
heterocyclic ring of 5 or 6 members;  
in the presence of a Lewis acid or fluoride reagent in an ether organic solvent at temperatures  
ranging from about -78°C to about 30°C to produce an alpha-sulfonyl carbonyl compound of  
formula V; any reactive substituent group(s) being protected during the reaction and removed  
thereafter; and further if desired isolating any chiral or stereoisomeric product as an  
individual isomer.